

Ethnobotanical Survey of Wild Edible Plants and Their Contribution for Food Security Used by Gumuz People in Kamash Woreda; Benishangul Gumuz Regional State; Ethiopia

Dessalegn Ayele Amente

Department of Biology, College of Natural and Computational Science, Assosa University, Assosa, Ethiopia

Email address:

dessalegnayele2011@gmail.com

To cite this article:

Dessalegn Ayele Amente. Ethnobotanical Survey of Wild Edible Plants and Their Contribution for Food Security Used by Gumuz People in Kamash Woreda; Benishangul Gumuz Regional State; Ethiopia. *Journal of Food and Nutrition Sciences*. Vol. 5, No. 6, 2017, pp. 217-224.

doi: 10.11648/j.jfns.20170506.12

Received: August 31, 2017; **Accepted:** September 23, 2017; **Published:** November 14, 2017

Abstract: The aims of this study was documenting and assessing the utilization of the food plants used by the Gumuz community in western Ethiopia. Informants were sampled from selected kebeles randomly. Semi-structured interview, questionnaires, focus group discussion and field observations were tools of data collection. Descriptive statistics, preference ranking, direct matrix ranking, and informant consensus were used to analyze the data. Total of 35 families, 49 genera with 60 species of wild edible plants were collected in the study area. Most of them are shrubs (36.67%) followed by trees and herbs with 28.33 % and 28.33% respectively. *Oxythenantra abyssinica* was the most preferred species. Wild edible plants are threatened due to various human and natural causes. Thus, public awareness and community based management is need to be encouraged by government and Non Government Organizations at all levels.

Keywords: Ethnobotany, Food Security, Wild Edible Plants

1. Introduction

Different People of the world feed upon various types of food depending on their culture. Thousands of species of wild fruits, leaves, seed, root and tuber are consumed as food worldwide. Wild edible plants (WEP) provide staple food for indigenous people, serve as complementary food for non-indigenous people and offer an alternative source of cash income for poor communities [1].

It is unclear how the current global food system will meet the growing demands of a population expected to reach 9 billion people by 2050 [2]. In the present context of climate change, ongoing loss of species and genetic diversity, soil degradation, increasing urbanization, social conflict and extreme poverty, there is an urgent need for collective action to address food and nutrition security at the global level [3].

WEPs play a great role in ensuring food security and improve the nutrition in the diets of many people in

developing countries [2]. Millions of people in many developing countries do not have enough food to meet their daily needs and a further more people are deficient in one or more micronutrients. Thus, in most cases rural communities depend on wild resources including WEPs to meet their food needs in periods of food crisis [4].

More recently, some ethno botanical studies have undertaken in some parts of Ethiopia. However, the majority of these studies have dealt with medicinal species and little emphasis has been paid to wild edible plants [4]. Since traditional knowledge on WEPs is being eroded through acculturation and the loss of plant biodiversity along with indigenous people and their cultural background, promoting research on wild food plants is crucial in order to safeguard this information for future [5]. WEP species are still largely ignored in land use planning and implementation, in economic development, and in biodiversity conservation endeavors. Therefore, this study was aimed to document such economic potentials for food security used by the Gumuz community of Kamash Woreda in western part of Ethiopia.

2. Materials and Methods

2.1. Description of the Study Area

The study was conducted during 2014-2016 in Kamash Woreda of Benishangul Gumuz Regional State which is located in the mid-western part of the Ethiopia. The Woreda is located on the western slopes of the Didessa River, with elevations ranging from approximately 2000 meters above sea level in the west to just under 1000 meters at the bottom of the Didessa valley. It is bordered by the Didessa River on the east which separates it from Yaso and Belo Jegonfo, by the Oromia Region on the south and west, and by Agalo Meti on the northwest.

The Central Statistical Agency national census reported a total population for this Woreda 17,883, of which 5,917 (33.09%) is urban dwellers and population density of 6.8 people per square kilometer which is less than the Zone average of 7.61. The two largest ethnic groups reported in Kamash Woreda were the Gumuz (81.4%), and the Oromo (17.4%); all other ethnic groups made up 1.2% of the population [6].

The average rain fall of the region is 1200mm in the area around Assosa and reduced towards north and west to about 800mm. Vegetation of the area is still nearly intact and has comparable vascular plant diversity to other floristic regions in Ethiopia.




Figure 1. Map of the study area.

2.2. Selection of Sites and Informants

A purposive sampling method was employed to select six (6) kebeles from the Woreda, Ten (10) randomly selected informants and six (6) purposively selected key informants from each kebele with total of 96 informants from the Woreda following [7].

2.3. Data Collection Methods

Semi-structured interview, direct matrix ranking, focus group discussions, and guided field walks with informants were employed following [7-9] and field observation was carried out with local field guide assistant who have the skill of local language.

2.4. Data Analysis

Descriptive statistics, preference ranking, paired comparison, direct matrix ranking, and informant consensus

were used to analyze the data.

3. Result and Discussion

3.1. Taxonomic Diversity of Wild Edible Plants

Total of 35 families, 49 genera with 60 species of WEPs were collected in the study area (figure 1). The family Fabaceae, Amaranthaceae and Acanthaceae were represented by the highest number of species (four species) or (6.67%) followed by Malvaceae, Asteraceae and Zingiberaceae that contributed three species (5%). Dioscoreaceae, Myrtaceae, Rubiaceae, Solanaceae, Tiliaceae, Rosaceae, Moraceae, Cucurbitaceae, Apocynaceae, and Vitaceae were the third in species diversity by having two species each (3%) and nineteen of the remaining families were represented by single species each (1.67%).

WEPs were also diversified in terms of their growth forms and accordingly most of them are shrubs accounting for

36.67% followed by trees and herbs with 28.33 % and 28.33% each. The least diversified wild edible plants in the area were found to be the climbers or lianas with 6.67% only (figure 2).

The result is different from the result obtained by [10] on

wild edible plants in which the trees were the mostly consumed growth forms followed by shrubs and this difference may come from the ecological variances and vegetation type of the two sites.




Figure 2. Taxonomic diversity wild edible plants in Kamash Woreda.




Figure 3. Wild edible plants diversity in their growth form.




Figure 2. Taxonomic diversity wild edible plants in Kamash Woreda.

3.2. Edible Parts Wild Edible Plants

From this study, 18 (30%) fruit and 14(23.33%) leaf were highly consumable parts of wild edible plants species by indigenous peoples of the study area. Tuber/Root 7(11.67%) and stem 5(8.33%) were next edible parts of the plants by the peoples. Seed 4(6.67%), Flower, stem & fruit, and fruit & seed each accounts for 3(5%), edible parts. The rest of parts like leaf and fruit, leaf and seed, and young bud and flower contributes 1(1.66%) each (figure 3).

This result similar with the study conducted on wild edible plants at Kara and Kwegu semi-pastoralist people in Lower Omo River Valley, Debub Omo Zone, SNNPR, of Ethiopia in which 23(58.97%) fruits, 13(33.33%) leaves, 3(7.69%) roots and one (2.56%) seed were edible parts of the plants in the area [10]. Not only this but other studies also supports this result in which fruits were reported as the most utilized plant parts out of the total parts used [11-12]. The same result was reported by Regassa in which fruit (40%) and fresh leaves (17.5%) followed by shoots (12.5%) were the most consumed parts of wild edible plants by Chellia peoples of Oromia Region, West-Central Ethiopia [13].

3.3. Contribution of Wild Edible Plants in Food Security

The general public consumes most of the WEPs as snacks, supplement or refreshments. So, most of the indigenous people of the area occasionally consider the WEPs as famine foods or foods in condition of starvation. According to the result from respondents, around 73.33% of the WEPs were used as supplementary food while the rest (26.67%) were used as food regularly or as meal (table 3). Similar studies

from other parts of Ethiopia; as well as from different parts of the world; reported that wild edible plants help to prevent starvation and sustain life during drought season and social unrest [14-17].

3.4. Preference Ranking of Wild Edible Plants by Informants

Species preference ranking of WEPs was conducted based on the informants' consensus approach to find out the relative importance of plants to the local community (Table 1). The most preferred species in increasing order were *Oxythenantra abyssinica*, *Abelmoschus esculentus*, *Corchorus olitorius*, *Dioscorea praehensilis*, *Annona senegalensis*, *Syzygium guineense* and *Luffa cylindrica*. The species preference is almost similar throughout the kebele and there is no significant difference among the six kebeles. This may be due to similarity in ethnic composition and sharing of the same culture of wild edible plant utilization as well living the same Woreda. *Oxythenantra abyssinica* was the most preferred edible plants, cited by most of respondents in all Kebele. Similarly, *Abelmoschus esculentus* and *Corchorus olitorius* were the best species cited by all respondents in all Kebeles as 2nd and 3rd in their preference while *Dioscorea praehensilis* is fourth. *Annona senegalensis* and *Syzygium guineense* were positioned at fifth and sixth rank of the selected wild edible plants of the area while *Luffa cylindrica* got the last position. Selection and ranking of the most preferred candidate of wild edible plant, based on their taste and utilization by the indigenous peoples of the area. The FGD participants also reflect the ranking.

Table 1. Results of preference ranking on seven most popular selected WEP species based their taste and repeated consumption by respondents of the study area (1=least, 2= less, 3=good, 4= very good and 5= excellent).

WEP species	Respondents							Total	Percent (%)	Rank
	R1	R2	R3	R4	R5	R6	R7			
<i>Abelmoschus esculentus</i>	4	5	4	5	4	4	5	31	17.61	2 nd
<i>Oxythenantra abyssinica</i>	5	5	4	5	5	4	5	33	18.75	1 st
<i>Dioscorea praehensilis</i>	4	3	3	4	3	5	4	26	14.77	4 th
<i>Corchorus olitorius</i>	4	4	5	4	3	4	3	27	15.34	3 rd
<i>Annona senegalensis</i>	4	3	4	3	2	3	4	23	13.07	5 th
<i>Syzygium guineense</i>	3	4	3	2	4	3	2	21	11.93	6 th
<i>Luffa cylindrica</i>	3	2	2	3	2	1	2	15	8.52	7 th
Total	27	26	25	26	23	24	25	176	99.99	

3.5. Disaster Coping Mechanisms During Food Shortage/Poverty

According to the responses from informants, using WEPs as sources food during famine is the first alternative to ensure food insecurity in the study area. It was stated as the wild foods from plants is not comparable to any other form mechanisms used to overcome food insecurity for indigenous peoples of the Gumuz ethnic groups in the study area (table 2). Hunting, collecting and sale of firewood, collection and sale of incense, government employee as day laborer & migration to other area are means of overcoming food insecurity.

The study conducted on Wild edible trees and shrubs in the semi-arid lowlands of Southern Ethiopia by Assefa and Abebe [19] supports the result of this study in that consumption of wild edible plant is the second mechanisms of coping with food insecurity in the area. The result of Guinand & Lemessa, [20]; Balemie & Kebebew, [4] on wild edible plants also revealed that, the consumption of wild food plants is ranked second as a coping mechanism for surviving during famine. The study conducted on Wild edible trees and shrubs in the semi-arid lowlands of Southern Ethiopia by Assefa and Abebe [19] supports the result of this study in that consumption of wild edible plant is the second mechanisms of coping with food insecurity in the area.

Table 2. Respondents' means of overcoming food insecurity (60 informants).

mechanism of handling food insecurity	Sample Kebeles						Total	Rank
	Hena	Daguba	Mirmita	Gilgila	Kobi	Jalo Leka		
Consumption of WEPs	4	4	5	3	4	3	23	1 st
Looking for gov't help	0	0	0	0	0	0	0	7 th
Collection and sale of firewood & charcoal	1	2	2	1	2	2	10	3 rd
Migration to other areas	1	1	0	0	1	0	3	6 th
Begging for food	0	0	0	0	0	0	0	7 th
Looking for employment as day-laborers	1	1	0	1	0	1	4	5 th
Hunting wild animals for meat	2	2	2	3	2	3	14	2 nd
Collection and sale of incense	1	0	1	2	1	1	6	4 th
Total	10	10	10	10	10	10	60	

3.6. Paired Comparison

For WEPs that were identified by the informants to be used as food, a paired comparison was made among five of them using ten informants to know their rank. Accordingly, *Oxythenantra abyssinica* stood first followed by *Ficus sur*

(table 3). This result indicates that *Oxythenantra abyssinica* is much favored over other plant species cited in the study area as food plants. Moreover, the result could be a testimony for the efficacy of these two plant species to use as the most preferable food plants at least in the study area.

Table 3. Result of Paired comparison of WEPs used by the people of the area (1 = Least; 2 = Good; 3 = Very good; 4 = Excellent).

Plants species	Respondents (R1-R10)										Total	Rank
	R1	R2	R3	R4	R5	R6	R7	R8	R9	R10		
<i>Ficus sur</i>	2	3	3	4	2	3	2	2	2	1	24	2 nd
<i>Cordia africana</i>	3	2	2	2	1	2	3	1	2	3	21	3 rd
<i>Syzygium guineense</i>	1	3	2	2	1	3	1	1	2	2	18	4 th
<i>Oxythenantra abyssinica</i>	4	4	3	4	4	3	4	4	4	3	37	1 st
<i>Amaranthus caudatus</i>	1	1	1	1	2	3	2	2	1	1	15	5 th

3.7. Use Diversity of Wild Edible Plants

Among the 60 WEPs documented in the study area, 48 plant species (80%) were reported to have multipurpose roles while 12 of them (20%) have only food role in the area (table 4). This finding shows that the local people harvest the wild edible plants of the area mostly for construction, firewood, and production of house hold equipment. The utilization of these plants for fire, construction and equipment is linked with the daily life activities of the community and the termination of their usage is impossible.

Table 4. Use diversity of wild edible plants in the study area.

Use	No of species	Percent (%)
Only food role	12	20
Food and firewood	10	16.6
Food and construction	13	21.7
Food and medicinal	7	11.7
Food and charcoal	5	8.3
Food and farm/house fence	9	15
Food and bee hive hanging	4	6.7

3.8. Threats to and Conservation Status

The ethno ecological knowledge on threats to WEPs and conservation concerns was also assessed. Like other plant species, WEPs are threatened due to various human activities and natural causes such as land use change (expansion of

agricultural lands), developmental activities (road construction and urbanization); habitat destruction (timber harvest, fuel wood collection, wild fire); drought, over-harvesting; and over-grazing were among the main factors that reduce the diversity and density of wild edible plants in the study area.

Although the potential impacts of climate change were also discussed; the respondents were reluctant to mention it as a major issue. It might be due to the fact that until recently the respondents did not experience and witness direct impacts of climate change on biodiversity.

To understand local people's perception on the factors more threatening WEP species with pair wise ranking of seven factors (agriculture, drought, and urbanization, fuel wood, wild fire, over grazing and selective cutting for construction) were conducted (table 5). The result indicated that agricultural activities and drought were ranked as first and second, while construction and over grazing was positioned at third and fourth stage followed by fuel wood, urbanization and wild fire which were ranked as fifth and sixth respectively. Similarly, the study conducted on Ethnobotanical study of wild edible plants in Derashe and Kucha Districts, Southern Ethiopia by Balemie and Kebebew [4] revealed that agricultural expansion as the principal threat to wild plant species.

Table 5. Direct matrix ranking results of six respondents on six factors perceived as threats to Wild edible plants (values: 1=least destructive, 6=most destructive).

Threatening factor	Respondents						Total	%age	Rank
	R1	R2	R3	R4	R5	R6			
Agriculture	6	5	5	4	6	4	30	18.75	1 st
Extended dry time	4	5	6	3	5	5	28	17.5	2 nd
Urbanization	3	2	3	1	3	4	16	10	6 th
Overgrazing	4	4	3	5	5	3	24	15	4 th
Construction	5	5	4	3	4	5	26	16.25	3 rd
Fuel wood	3	4	3	4	2	4	20	12.5	5 th
Wild fire	4	2	3	2	2	3	16	10	6 th

4. Conclusion

The result of the study revealed that knowledge about the edibility, habitat distribution, harvesting time and uses of most wild edible plant species is still preserved among the study communities. The preservation of this knowledge appears to be the result of continued reliance of local communities on the wild edible plants and it's passed from elders to younger's orally. Analysis of the results showed that in all study kebeles, most of the edible plants were used mainly by all communities during normal and shortage of food. Utility of the wild edible plants especially by children of the community members indicated the maintenance of indigenous knowledge associated with the plant species in the area. But the diversity of wild edible plants in the area is declined gradually by different factors leading to the fading away of this indigenous knowledge associated with the plants of the area. The results also showed that many wild edible species are under growing pressures from various anthropogenic factors. Thus, public awareness and community based management need to be encouraged by government and non government organizations at all levels with urgent collection of Germplasm by the professionals. The findings suggest further investigation into nutritional profiles and processing methods of all the species reported and study of the pharmacological properties for the nutraceutical species since they are also used for medicinal applications.

Recommendation

Based on the results of the study, the following recommendations are forwarded:

1. Local communities should be encouraged to cultivate wild food plants on their own land. Provision for technical as well as financial assistance should be made in this regard.
2. Raising awareness of the young generation to avoid negative impacts on wild edible plants and associated knowledge. So that the plants and the TK would be preserved.
3. The indigenous knowledge and skill of using wild edible plants must be encouraged and protected.
4. Establishing conservation measures and strategies to ensure the sustainability of multipurpose and widely used wild edible plants is needed as most them are obtained from the wild. This can be achieved by:
 - (a). Encouraging people to grow wild edible plants in the home gardens, mixing with crops in farmlands and live fences.
 - (b). Promoting the establishment of local botanical garden starting at least at the Woreda level.
5. Since some of the highly valued wild edible plants are being over exploited due to their use for food purposes, specific wild food plant conservation strategy should be formulated and implemented for long term management of plants in the area.
6. A long term research and monitoring on ethnobotany and its practices on other wealth of useful plants from the forests and surroundings is imperative.
7. The findings suggest further investigation into nutritional profiles and processing methods of all the species reported and study of the pharmacological properties for the nutraceutical species since they are also used for medicinal applications.

Appendix

Table 6. Local name, scientific name, parts used and habit of Collected WEP Species in the Study Area.

No	Local name(Gumuz)	Family	Scientific name	Parts used	Habit
1	Andegila	Bignoniaceae	<i>Stereospermum kunthianum</i> Cham.	Young stem	Shrub
2	Agidema	Asteraceae	<i>Vernonia theophrastifolia</i> schweinf.ex Oliv& Hiern	Leaf	Shrub
3	Ajanzibil	Zingiberaceae	<i>Zingiber officinale</i> Roscoe	Root	Herb
4	Akala	Fabaceae	<i>Vigna subterranea</i> (L.) Verdc.	Seed	Shrub
5	Akechechile	Amaranthaceae	<i>Amaranthus caudatus</i> L.	Cooked Leaf	Herb
6	Ambershuwa	Amaranthaceae	<i>Celosia trigyna</i> L.	Cooked Leaf	Herb
7	Andedha	Malvaceae	<i>Abelmoschus esculentus</i> (L.) Moench.	Fruit	Shrub
8	Andeha	Malvaceae	<i>Abelmoschus ficulneus</i> (L.) weight & Arn	Leaf & fruit	Shrub
9	Anjidema	Asteraceae	<i>Vernonia theophrastifolia</i> Schweif.ex Oliv.& Heim	Leaf	Shrub

No	Local name(Gumuz)	Family	Scientific name	Parts used	Habit
10	Antsiqina	Vitaceae	<i>Ampelocissus schiperiana</i> (L.) weight & Arn	Young stem	Shrub
11	Antsitsa	Flacourtiaceae	<i>Oncoba spinosa</i> Forssk.	fruit & young stem	Shrub
12	Baga	Brassicaceae	<i>Brassica carinata</i> A. Br.	Leaf	Shrub
13	Bakeya	Rubiaceae	<i>sarcocephalus latifolius</i> (J. E. Smith) E. A	fruit	Tree
14	Bambeya	Convolvulaceae	<i>Ipomea batatas</i> L.	Root	Liana
15	Bambuta	Annonaceae	<i>Annona senegalensis</i> Pers.	fruit	Tree
16	Banja	Boraginaceae	<i>Cordia africana</i> Lam.	fruit	Tree
17	Bedaha	Cucurbitaceae	<i>Luffa cylindrica</i> (L.) M. J. Roem.	Cooked leaf	Herb
18	Beewa	Fabaceae	<i>Lonchocarpus laxiflorus</i> Guill. & Perr.	Grinned flower	Tree
19	Bela	Portulacaceae	<i>Portulaca oleracea</i>	Leave & seed	Shrub
20	Bishela	Apiaceae	<i>Anethum graveolens</i> L.	Flower	Shrub
21	Boha	Acanthaceae	<i>Acanthus pubescens</i> (Oliv.) Engl.	Flower	shrub
22	Chichariya	Amaranthaceae	<i>Amaranthus spinosus</i> L.	Leave	Shrub
23	Darguwa	Acanthaceae	<i>Asystasia gangetica</i> (L.) T.Andress.	Cooked leaf	Herb
24	Diwa	Myrtaceae	<i>Syzygium guineense</i> (Willd.) DC. subsp guineense	fruit	Tree
25	Doga	Fabaceae	<i>Tamarindus indica</i> L.	Ripped seed & fruit	Tree
26	Ebosiya	Solanaceae	<i>Solanum alatum</i> Moench.	Leaf	Herb
27	Eca	Dioscoreaceae	<i>Dioscorea praeheensis</i> Benth.	root	Liana
28	Echa	Moraceae	<i>Ficus sur</i> Forssk.	Fruit	Tree
29	Ejega	Malvaceae	<i>Hibiscus sabdariffa</i> L.	Fruit & seed	Shrub
30	Ejimbeya	Anacardiaceae	<i>Ozoroa pulcherrima</i> (Schweinf.) R.& A. Fernand	Stem	Herb
31	Ejisiya	Solanaceae	<i>Physalis peruviana</i> L.	Young bud & Flower	Tree
32	Ekicanqila	Amaranthaceae	<i>Amaranthus hybridus</i>	Cooked leaf	Herb
33	Elangiya	Acanthaceae	<i>Justicia ladanoides</i> Lam.	Cooked Leaf	Herb
34	Ewa	Vitaceae	<i>Cissus populnea</i> Guill. & Perr.	Leaves	Shrub
35	Feya	Olaceae	<i>Ximenia Americana</i> L.	Fruit	Tree
36	Gediya	Tiliaceae	<i>Grewia mollis</i> A. Juss	Fruit	Tree
37	Hafa	Combretaceae	<i>Combretum collinum</i> Fresen.	Seed	Tree
38	Hangua	Rhamnaceae	<i>Ziziphus abyssinica</i> Hochst ex A. Rich	Fruit	Tree
39	Heca /qilxu	Moraceae	<i>Ficus vasta</i> Forssk.	Fruit	Tree
40	Heya	Olaceae	<i>Ximenia americana</i> L.	fruit	Herb
41	Hojiya /enta	Poaceae	<i>Oxythenantra abyssinica</i> (A. Rich) Munro	Young stem	Herb
42	Huya	Apocynaceae	<i>Saba comorensis</i>	Fruit & seed	Liana
43	Igidimba	Asteraceae	<i>Vernonia auriculifera</i> Hiern.	Cooked leaf	Shrub
44	kompha	Araceae	<i>Colocasia esculenta</i> (L.) Schott.	Root	Herb
45	Kota	Rubiaceae	<i>Gardenia ternifolia</i> Schum. & Thonn.	fruit	Tree
46	Lalqa	Tiliaceae	<i>Corchorus olitorius</i> L.	Leaf	Shrub
47	Mecha	Fabaceae	<i>Piliostigma thonningii</i>	fruit	Tree
48	Molowa/Metiya	Arecaceae	<i>Phoenix reclinata</i> Jacq.	Young stem & fruit	Herb
49	Ola	Zingiberaceae	<i>Aframomum alboviolaceum</i> (Ridl.) K. schum	Raw root	Herb
50	Muga	Acanthaceae	<i>Justicia schimperiana</i> (Hochst.ex Nees) T. Anders	Flower	Shrub
51	Qoqora	Verbenaceae	<i>Vitex doniana</i> Sweet.	Ripped fruit	Tree
52	Qosha	Euphorbiaceae	<i>Ricinus communis</i> L.	Seed	Shrub
53	Siwa	Apocynaceae	<i>Carissa spinarum</i> L.	Fruit	Shrub
54	Shawa	Myrtaceae	<i>guineense</i> (Willd.) DC. supsp <i>afromontanum</i>	fruit	tree
55	Waga	Rosaceae	<i>Rubus steudneri</i> Shweinf.	Fruit	Shrub
56	Xasha	Celastraceae	<i>Maytenus senegalensis</i> (Lam.) Exell.	Burned stem as salt	Herb
57	Yampedema	Zingiberaceae	<i>Costus spectabilis</i> (Fenzl) K. schum	Root	Shrub
58	Yechiwa	Dioscoreaceae	<i>Dioscorea bulbifera</i>	Root	Liana
59	Sinada	Rosaceae	<i>Rubus apetalus</i> Poir.	fruit	Shrub
60	Equmetsa	Cucurbitaceae	<i>Peponium vogelii</i> (Hook.f.) Engl.	Fruit	Herb

References

- [1] Ju, Y., Zhuo, J., Liu B. and Long, C. Eating from the wild: diversity of wild edible plants used by Tibetans in Shangri-la region, Yunnan, China. *Journal of Ethnobiology and Ethnomedicine* (2013); 9:28
- [2] Food and Agricultural Organization. Forests for Improved Nutrition and Food Security (2010).
- [3] Hunter, D. and Fanzo, J. Agricultural biodiversity, diverse diets and improving nutrition (2013). London, Earth scan publishers.
- [4] Balemie K and Kebebew F. Ethnobotanical study of wild edible plants in Derashe and Kucha Districts, South Ethiopia. *J Ethnobiol and Ethnomed.* (2006). 2:53
- [5] Lulekal E, Asfaw Z, Kelbessa E and Damme, V. P. Wild edible plants in Ethiopia: a review on their potential to combat food insecurity (2011).
- [6] Central Statistical Authority. National Population and Housing Census of Ethiopia. Addis Ababa (2007).
- [7] Martin GJ. *Ethnobotany: A methods manual*. Chapman and Hall, London; (1995).87-105
- [8] Cotton, C. M. *Ethnobotany: Principles and Applications*. John Wiley and Sons, New York, (1996). 412.

- [9] Alexiades, M. N. Collecting ethnobotanical data: An introduction to basic concepts and techniques. *Selected Guidelines for Ethnobotanical Research: A Field Manual*. The New York Botanical Garden, Bronx, New York (1996). 53-94.
- [10] Teklehaymanot T. and Giday M. Ethnobotanical study of wild edible plants of Kara and Kewego semipastoralist people in Lower Omo River valley, Debub Omo Zone, SNNPR, Ethiopia. *J Ethnobioland Ethnomed* (2010). 6: 23.
- [11] Tamene B, Bekele T, and Kelbessa E. An Ethnobotanical Study of the Semi-Wetland Vegetation of Cheffa. Addis Ababa University Press, Addis Ababa, Ethiopia, 2000, <http://www.phe-ethiopia.org/pdf/Hujub%20Sacred%20Forests.pdf>.
- [12] Wondmu T, Asfaw Z and Kelbessa E. Ethnobotanical study of foodplants around deeraa town arsi, Ethiopia. *SINET: Ethiopian Journal of Science*. (2006). 29(1):71-80.
- [13] Regassa T, Kelbessa E and Asfaw Z. Ethnobotany of Wild and Semi- Wild Edible Plants of Chelia District, West-Central Ethiopia; *Sci. Technol. Arts Res. J.*, (2014).3(4): 122-134.
- [14] Zemedu A. Indigenous African food crops and useful plants: Survey of indigenous food crops, their preparations and home gardens Nairobi: The United Nation University Institute for Natural Resources in Africa. (1997).
- [15] Awas T. Plant diversity in western Ethiopia: Ecology, Ethnobotany and Conservation (1997).
- [16] Neudeck L, Avelino L, Bareetseng P, Barbara Ngwenya N, Teketay D and Moseki R. *Journal of Ethnobotany Research & Applications* (2012).10:449-462.
- [17] Cunningham, A. B. Applied Ethnobotany: People, Wild Plant Use and Conservation. People and Plant Conservation manual. Earth scans Publication Ltd. (2001). 145
- [18] Grivetti, L. (2001). Reading 8: Edible Wild Plants, Part 1, issued on 15/10/2016. <http://www.ucdavis.-edu\Grivetti.htm>
- [19] Assefa A & Abebe T. Wild Edible Trees and Shrubs in the Semi-arid Lowlands of Southern Ethiopia. *Journal of Science & Development* (2011). 1: 5-19.
- [20] Guinand, Y. and Dechassa L. Wild-food plants in Southern Ethiopia: Reflections on the role of famine-foods at a time of drought. UNDP-EUE. Field mission report, Addis Ababa. (2000).